

Sudden Deafness

Description

Sudden Sensorineural Hearing Loss (SSHL), or sudden deafness, is a rapid loss of hearing. SSHL can happen to a person all at once or over a period of up to 3 days. It should be considered a medical emergency. A person who experiences SSHL should visit a doctor immediately.

A doctor can determine whether a person has experienced SSHL by conducting a normal hearing test. If a loss of at least 30 decibels in three connected frequencies is discovered, it is diagnosed as SSHL. A decibel is a measure of sound. A decibel level of 30 is half as loud as a normal conversation. A frequency is another way of measuring sound. Frequencies measure sound waves and help to determine what makes one sound different from another sound.

Hearing loss affects only one ear in 9 out of 10 people who experience SSHL. Many people notice it when they wake up in the morning. Others first notice it when they try to use the deafened ear, such as when they make a phone call. Still others notice a loud, alarming "pop" just before their hearing disappears. People with SSHL often experience dizziness or a ringing in their ears (**tinnitus**), or both.

Some patients recover completely without medical intervention, often within the first 3 days. This is called a **spontaneous recovery**. Others get better slowly over a 1 or 2 week period. Although a good to excellent recovery is likely, 15 percent of those with SSHL experience a hearing

loss that gets worse over time.

Approximately 4,000 new cases of SSHL occur each year in the United States. It can affect anyone, but for unknown reasons it happens most often to people between the ages of 30 and 60.

Causes/Diagnosis

Though there are more than 100 possible causes of sudden deafness, it is rare for a specific cause to be precisely identified. Only 10 to 15 percent of patients with SSHL know what caused their loss. Normally, diagnosis is based on the patient's medical history. Possible causes include the following:

- Infectious diseases.
- Trauma, such as a head injury.
- Abnormal tissue growth.
- Immunologic diseases such as Cogan's syndrome.
- Toxic causes, such as snake bites.
- Ototoxic drugs (drugs that harm the ear).
- Circulatory problems.
- Neurologic causes such as multiple sclerosis.
- Relation to disorders such as Ménière's disease.

Treatment

People who experience SSHL should see a physician immediately. Doctors believe that finding medical help fast

increases the chances for recovery. Several treatments are used for SSHL, but researchers are not yet certain which is the best for any one cause. If a specific cause is identified, a doctor may prescribe antibiotics for the patient. Or, a doctor may advise a patient to stop taking any medicine that can irritate or damage the ear.

The most common therapy for SSHL, especially in cases with an unknown cause, is treatment with steroids. Steroids are used to treat many different disorders and usually work to reduce inflammation, decrease swelling, and help the body fight illness. Steroid treatment helps some SSHL patients who also have conditions that affect the immune system, which is the body's defense against disease.

Another common method that may help some patients is a diet low in salt. Researchers believe that this method aids people with SSHL who also have Ménière's disease, a hearing and balance disorder.

Research

Two factors that help hearing function properly are good air and blood flow inside the ear. Many researchers now think that SSHL happens when important parts of the inner ear do not receive enough oxygen. A common treatment for this possible cause is called carbogen inhalation. Carbogen is a mixture of oxygen and carbon dioxide that seems to help air and blood flow better inside the ear. Like steroid therapy, carbogen inhalation does not help every patient, but some SSHL patients taking carbogen have recovered over a period of time.

Additional Resources

American Academy of Otolaryngology-Head and Neck Surgery (AAO-HNS)

One Prince Street

Alexandria, VA 22314-3357

Voice (703) 836-4444

TTY: (703) 519-1585

Fax: (703) 683-5100

E-mail: webmaster@entnet.org

Internet: www.entnet.org

Self Help for Hard of Hearing People, Inc. (SHHH)

7910 Woodmont Avenue

Suite 1200

Bethesda, MD 20814

Voice: (301) 657-2248

TTY: (301) 657-2249

Fax: (301) 913-9413

E-mail: national@shhh.org

Alexander Graham Bell Association for the Deaf and Hard of Hearing (A.G. Bell)

3417 Volta Place, NW

Washington, DC 20007

Voice/TTY: (202) 337-5220

Toll Free: (800) HEAR-KID

Fax: (202) 337-8314

E-mail: info@agbell.org

Internet: www.agbell.org

Association of Late Deafened Adults (ALDA)

1131 Lake Street
Box 204
Oak Park, IL 60301
Toll Free Voice/Fax: (877) 907-1738
TTY: (708) 358-0135
E-mail: info@alda.org
Internet: www.alda.org

CHID database search

CHID is a database produced by health-related agencies of the Federal Government. This database provides titles, abstracts, and availability information for health information and health education resources. The value of this database is that it lists a wealth of health promotion and education materials and program descriptions that are not indexed elsewhere. Search the database using "sudden deafness" to view citations to journal articles, educational materials, and books on this topic.

PubMed database search

PubMed is a database developed by the National Library of Medicine in conjunction with publishers of biomedical literature as a search tool for accessing literature citations and linking to full-text journals at web sites of participating publishers. Search the database using "sudden deafness" for medical journal articles.

NIH Pub. No. 00-4757
February 2000
Updated March 2003